A Note on Skew Characters of Symmetric Groups

Jay Taylor

Abstract. In previous work Regev used part of the representation theory of Lie superalgebras to compute the values of a character of the symmetric group whose decomposition into irreducible constituents is described by semistandard (k,ℓ)-tableaux. In this short note we give a new proof of Regev's result using skew characters.

arXiv:1603.03914v1 [math.RT] 12 Mar 2016

1. Introduction

1.1. For any partition $\alpha \in \mathsf{P}(n)$ of an integer $n \geqslant 0$ we have a corresponding irreducible character χ_α of the symmetric group $\mathfrak{S}_n$; we assume this labelling is as in [Mac95, 7.4]. In [Reg13] Regev observed that the values of the character $\Gamma_n = \sum_{a=0}^{n} \chi_{(a,1^{n-a})}$ obtained by summing over all hook partitions were particularly simple. Specifically if $\nu \in \mathsf{P}_r(n)$ is a partition of length r then we have

$$\Gamma_n(\nu) = \begin{cases} 2^{r-1} & \text{if all parts of } \nu \text{ are odd,} \\ 0 & \text{otherwise.} \end{cases}$$

Here we write $\Gamma_n(\nu)$ for the value of Γ_n at an element of cycle type ν.

1.2. To prove this result Regev considered a more general but related problem which we now recall. For any integers $k,\ell \geqslant 0$ and any partition $\alpha \in \mathsf{P}(n)$ we denote by $s_{k,\ell}(\alpha)$ the number of all semistandard (k,ℓ)-tableaux of shape α, see 2.6 for the definition. Motivated by the representation theory of Lie superalgebras Regev considered the following character of $\mathfrak{S}_n$

$$\Lambda_n^{k,\ell} = \sum_{\alpha \in \mathsf{P}(n)} s_{k,\ell}(\alpha)\chi_\alpha.$$

The main result of [Reg13] is the following.

Theorem 1.3 (Regev). *If $\nu = (\nu_1, \dots, \nu_r) \in \mathsf{P}_r(n)$ is a partition of length r then*

$$\Lambda_n^{k,\ell}(\nu) = \prod_{i=1}^{r} (k + (-1)^{\nu_i - 1}\ell).$$

Remark 1.4. Note that $s_{k,\ell}(\alpha) \neq 0$ if and only if α is contained in the (k,ℓ)-hook, as defined in [BR87, 2.3]. In particular, we have $\Lambda_n^{k,\ell}$ is the same as the character $\chi_{\varphi^*_{(k,\ell),n}}$ defined in [Reg13].

2010 Mathematics Subject Classification. Primary 20C30; Secondary 20C15.
Keywords and phrases. Symmetric groups, representation theory, skew characters, Murnaghan–Nakayama formula..

1.5. Using formulas for the coefficients $s_{k,\ell}(\alpha)$ obtained in [BR87] Regev deduces that $\Lambda_n^{1,1} = 2\Gamma_n$ from which the statement of 1.1 follows immediately. To prove Theorem 1.3 Regev used results of Berele–Regev on representations of Lie superalgebras [BR87]. However, Regev asks whether this result can be proven entirely in the setting of the symmetric group. The purpose of this note is to provide such a proof. Our proof is based on a description of the character $\Lambda_n^{k,\ell}$ as a sum of skew characters. With this we can use the Murnaghan–Nakayama formula to compute the values of $\Lambda_n^{k,\ell}$ and thus prove Theorem 1.3. As a closing remark we use our description in terms of skew characters to show that $\Lambda_n^{1,1} = 2\Gamma_n$ using Pieri's rule.

Remark 1.6. The idea we use in this paper was prompted by a recent question of Marcel Novaes on MathOverflow [Nov16], which is where we also first learned of Regev's work.

Acknowledgments: The author gratefully acknowledges the financial support of an INdAM Marie-Curie Fellowship and grants CPDA125818/12 and 60A01-4222/15 of the University of Padova. Furthermore we thank Chris Bowman for sharing his thoughts on an earlier version of this article.

2. The Result

2.1. Let $\mathbb{N} = \{1, 2, \dots\}$ be the natural numbers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Throughout we use the term *diagram* to mean a subset of $\mathbb{N}^2$. The notion of connected diagram and connected components of a diagram have their usual natural meanings, see [Mac95, I, §1] for details. A diagram T will be called a *horizontal line*, resp., *vertical line*, if for any $(i, j), (i', j') \in T$ we have $i = i'$, resp., $j = j'$.

2.2. For any $k \in \mathbb{N}_0$ we denote by C_k the set of all compositions $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{N}_0^k$ of length k; we call α_i a *part* of α. For such a composition we denote by $|\alpha|$ the sum $\alpha_1 + \cdots + \alpha_k$ and by α° the composition obtained from α by removing all parts equal to 0 but maintaining the original order. If $n \in \mathbb{N}_0$ then we denote by $\mathsf{P}_k(n)$ the set of all $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathsf{C}_k$ such that $\alpha_1 \geqslant \cdots \geqslant \alpha_k > 0$ and $|\alpha| = n$, which are the partitions of n of length k. Moreover we denote by $\mathsf{P}(n)$ the set $\bigcup_{k\in\mathbb{N}} \mathsf{P}_k(n)$ of all partitions of n. To each partition $\alpha \in \mathsf{P}(n)$ we have a corresponding diagram $T_\alpha = \{(i, j) \mid 1 \leqslant j \leqslant \alpha_i\}$ called the *Young diagram* of α. A diagram S is then called a *skew diagram* if $S = T_\alpha \setminus T_\beta$ for some Young diagrams $T_\beta \subseteq T_\alpha$. We recall that if S is a skew diagram with $|S| = n$ then we have a corresponding character ψ_S of $\mathfrak{S}_n$ called a *skew character*, c.f., [Mac95, §7, Example 3]. The following property of these characters is well known.

Lemma 2.3 (see [Mac95, 5.7]). *If S and S' are skew diagrams with the same connected components then $\psi_S = \psi_{S'}$.*

2.4. For any $k, \ell, n \in \mathbb{N}_0$ we denote by $\mathsf{B}_{k,\ell}(n) \subseteq \mathsf{C}_k \times \mathsf{C}_\ell$ the set of all pairs $(\lambda \mid \mu)$ of compositions such that $|\lambda| + |\mu| = n$; we call these *bicompositions* of n. Now for each bicomposition $(\lambda \mid \mu) \in \mathsf{B}_{k,\ell}(n)$ we denote by $S_{(\lambda|\mu)}$ some (any) skew diagram whose connected components $H_1, \dots, H_r, V_1, \dots, V_s \subseteq S_{(\lambda;\mu)}$ are such that H_i is a horizontal line, resp., V_j is a vertical line, and $(|H_1|, \dots, |H_r|) = \lambda^\circ$, resp., $(|V_1|, \dots, |V_s|) = \mu^\circ$. It is easy to see that such a diagram exists. By Lemma 2.3 we then get a well defined character $\psi_{(\lambda|\mu)} := \psi_{S_{(\lambda|\mu)}}$ of $\mathfrak{S}_n$.

Example 2.5. Consider the bicomposition $(4, 0, 5; 2, 3) \in \mathsf{B}_{3,2}(14)$ then an example of a corresponding skew diagram $S_{(4,0,5;2,3)}$ is given by



2.6. We now prove Theorem 1.3 but before proceeding we recall some definitions from [BR87, 2.1]. Specifically, let $D = \{1, \dots, k, 1', \dots, \ell'\}$ be a totally ordered set with $1 < \cdots < k < 1' < \cdots < \ell'$. If $\alpha \in \mathsf{P}(n)$ is a partition and $(\lambda \mid \mu) \in \mathsf{B}_{k,\ell}(n)$ is a bicomposition then we say a function $f : T_\alpha \to D$ is a (k,ℓ)-tableau of shape α and weight $(\lambda \mid \mu)$ if $\lambda_i = |\{x \in T_\alpha \mid f(x) = i\}|$ for any $1 \leqslant i \leqslant k$ and $\mu_j = |\{x \in T_\alpha \mid f(x) = j'\}|$ for any $1 \leqslant j \leqslant \ell$. As in [BR87, 2.1] we say f is *semistandard* if $T_f = f^{-1}(\{1, \dots, k\})$ is a Young tableau whose rows are weakly increasing and whose columns are strictly increasing and $T_\alpha \setminus T_f$ is a skew tableau whose columns are weakly increasing and whose rows are strictly increasing. If $s_{(\lambda|\mu)}(\alpha)$ is the number of semistandard (k,ℓ)-tableaux of shape α and weight $(\lambda \mid \mu)$ then $s_{k,\ell}(\alpha) := \sum_{(\lambda|\mu)} s_{(\lambda|\mu)}(\alpha)$ is the number of all semistandard (k,ℓ)-tableaux of shape α.

Lemma 2.7. *For any $k, \ell, n \in \mathbb{N}_0$ we have*

$$\Lambda_n^{k,\ell} = \sum_{(\lambda|\mu) \in \mathsf{B}_{k,\ell}(n)} \psi_{(\lambda|\mu)}.$$

Proof. It follows from [Mac95, 3.4, 5.1, 5.4, 7.3] that for any bicomposition $(\lambda \mid \mu) \in \mathsf{B}_{k,\ell}(n)$ we have

$$\psi_{(\lambda|\mu)} = \mathrm{Ind}_{\mathfrak{S}_\lambda \times \mathfrak{S}_\mu}^{\mathfrak{S}_n}(\chi_{(\lambda_1)} \boxtimes \cdots \boxtimes \chi_{(\lambda_k)} \boxtimes \chi_{(1^{\mu_1})} \boxtimes \cdots \boxtimes \chi_{(1^{\mu_\ell})}),$$

where $\mathfrak{S}_\lambda \times \mathfrak{S}_\mu$ is the Young subgroup determined by the parts of λ and μ. By [BR87, Lemma 3.23] we have the decomposition of this character into irreducibles is given by

$$\psi_{(\lambda|\mu)} = \sum_{\alpha \in \mathsf{P}(n)} s_{(\lambda|\mu)}(\alpha) \chi_\alpha.$$

Note that when $\ell = 0$ this statement is just Young's rule and as in [BR87] the general case can be proved easily by induction on ℓ using the definition of (k,ℓ)-tableaux. With this we obtain the desired statement

$$\sum_{(\lambda|\mu) \in \mathsf{B}_{k,\ell}(n)} \psi_{(\lambda|\mu)} = \sum_{(\lambda|\mu) \in \mathsf{B}_{k,\ell}(n)} \sum_{\alpha \in \mathsf{P}(n)} s_{(\lambda|\mu)}(\alpha) \chi_\alpha = \sum_{\alpha \in \mathsf{P}(n)} s_{k,\ell}(\alpha) \chi_\alpha. \qquad \blacksquare$$

Proof (of Theorem 1.3). Choose a part a of ν and let $\hat{\nu} \in \mathsf{P}(n-a)$ be the partition obtained by removing the part a from ν but maintaining the original order. If $\lambda \in \mathsf{C}_k$ is a composition such that $\lambda_i \geqslant a$ then we denote by $\lambda \downarrow_i a \in \mathsf{C}_k$ the composition obtained by replacing λ_i with $\lambda_i - a$. By the Murnaghan–Nakayama formula for skew characters, see [JK81, 2.4.15], we have that

$$\psi_{(\lambda|\mu)}(\nu) = \sum_{\lambda_i \geqslant a} \psi_{(\lambda \downarrow_i a|\mu)}(\hat{\nu}) + \sum_{\mu_j \geqslant a} (-1)^{a-1} \psi_{(\lambda|\mu \downarrow_j a)}(\hat{\nu})$$

where the first, resp., second, sum is over all $1 \leqslant i \leqslant k$, resp., $1 \leqslant j \leqslant \ell$, such that $\lambda_i \geqslant a$, resp., $\mu_j \geqslant a$. Indeed, the connected component of the skew diagram $S_{(\lambda|\mu)}$ labelled by λ_i, resp., μ_j, has an a-hook of leg length 0, resp., $(-1)^{a-1}$, if and only if $\lambda_i \geqslant a$, resp., $\mu_j \geqslant a$. Now clearly every bicomposition $(\lambda' \mid \mu') \in \mathsf{B}_{k,\ell}(n-a)$ arises from exactly $k+\ell$ bicompositions $(\lambda \mid \mu) \in$ via the process $\downarrow_i a$ and so by Lemma 2.7 we have

$$\Lambda_n^{k,\ell}(\nu) = \sum_{(\lambda|\mu)\in \mathsf{B}_{k,\ell}(n)} \psi_{(\lambda|\mu)}(\nu) = (k + (-1)^{a-1}\ell)\Lambda_{n-a}^{k,\ell}(\hat{\nu}).$$

An easy induction argument completes the proof. ■

Remark 2.8. Recall that the decomposition of $\psi_{(\lambda|\mu)}$ into irreducible characters is given by the Littlewood–Richardson coefficients, see [Mac95, 5.3]. If $k = \ell = 1$ then a simple application of Pieri's rule shows that

$$\psi_{(a|n-a)} = \begin{cases} \chi_{(1^n)} & \text{if } a = 0, \\ \chi_{(a,1^{n-a})} + \chi_{(a+1,1^{n-a-1})} & \text{if } 0 < a < n, \\ \chi_{(n)} & \text{if } a = n. \end{cases}$$

This gives an alternative way to see that $\Lambda_n^{1,1} = 2\Gamma_n$.

References

[BR87] A. Berele and A. Regev, *Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras*, Adv. in Math. **64** (1987), no. 2, 118–175.

[JK81] G. James and A. Kerber, *The representation theory of the symmetric group*, vol. 16, Encyclopedia of Mathematics and its Applications, With a foreword by P. M. Cohn, With an introduction by Gilbert de B. Robinson, Addison-Wesley Publishing Co., Reading, Mass., 1981, xxviii+510.

[Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Second, Oxford Mathematical Monographs, With contributions by A. Zelevinsky, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995, x+475.

[Nov16] M. Novaes, *Sum of skew characters over hooks and "odd" partitions*, MathOverflow, Mar. 2016, URL: `http://mathoverfl`

[Reg13] A. Regev, *Lie superalgebras and some characters of S_n*, Israel J. Math. **195** (2013), no. 1, 31–35.

Dipartimento di Matematica, Universitá di Padova, Via Trieste 63, 35121 Padova, Italy.
Email: `taylor@math.unipd.it`